Examining Colorectal and Cervical Cancer Care in Appalachia

A review of barriers and interventions to cancer screening, genetic services, and continuity of care

By Richard Ingram, MD; Molly Black; Susan Garwood, MD; Aasems Jacob, MD, FACP, FAPCR; Jeffrey Kendall, PsyD, LP; Richard Martin III, MD, MPH; Martin Palmeri, MD, MBA; Sashi Naidu, MD; Nicole Strout, DPT, CLT-LANA, FAPTA; Melissa Thomas, PhD, MSPH, MSA, MCHES, C-CHW; David Switzer, MD, FAAFP; Serena Phillips, DrPH, RN; Allison Harvey, MPH, CHES; Elana Plotkin, CMP-HC; Savannah Dodson, MPH; Stephanie Helbling, MPH, MCHES; and Leigh Boehmer, PharmD, BCOP

Appalachia encompasses 13 states, spanning 206,000 square miles.¹ It is home to nearly 26 million people. A fourth of Appalachia's 423 counties are rural, characterized by generational poverty and a lack of key resources, such as adequate healthcare. This is evidenced in an overall Appalachian cancer mortality rate that runs 10 percent higher than the national average.² Appalachia's most rural populations experience cancer death rates more than 15 percent higher than the rest of the nation. In central Appalachia, the mortality rate is 32 percent higher than the rest of the U.S. In fact, Appalachian Kentucky has the highest rates of cancer burden, exceeding the national average by 35 percent.

Lung, cervical, and colorectal cancer incidence and mortality are higher in the Appalachian areas of Ohio, Kentucky, Pennsylvania, Virginia, and West Virginia than any other parts of the U.S.³ In addition, there is a general lack of genetics professionals in Appalachia, and services tend to be clustered around major cities resulting in barriers of distance, transportation, and time.⁴⁻⁶ Other barriers to genetic screening and counseling include lack of awareness about genetic testing and potential benefits, low demand, perception of high cost, low prioritization, lack of physician recommendation, and lack of insurance coverage.^{4,5}

With multiple providers delivering care to patients with cancer, there are increased opportunities for gaps in communication and role confusion among primary care providers (PCPs) and cancer specialists. The transition from active treatment to survivorship care is a critical piece to patients' long-term health outcomes. Barriers faced by patients during this transition of care may include fragmented care, poor communication (especially among providers and patients), and a lack of understanding of future risks or a follow-up plan.⁷

The Association of Community Cancer Centers (ACCC) has joined with state oncology societies from throughout the region to create the Appalachian Community Cancer Alliance (the "Alliance"). The Alliance seeks to provide residents of Appalachia and beyond with interdisciplinary, patient-centered approaches to cancer care from prevention through survivorship, with an emphasis on enhanced quality-of-life. A landscape analysis was conducted to inform the Alliance's strategic planning by providing an overview of current locoregional activities, barriers, and interventions around colorectal and cervical cancer screenings, genetics services, and continuity of care along the cancer continuum.

Barriers

Barriers to care, and especially cancer screenings, are well documented in all patient populations. There are often multiple, confounding factors, for example, patient fear, community distrust, cost, and lack of transportation. Some barriers are out of patient control, such as distance to a healthcare facility, inconvenient clinic hours, or limited capacity. Reducing such structural barriers leads to increased access to cancer screenings.

ACCC's literature review identified the following barriers to screenings and related services.

Cervical Cancer

Incidence and mortality rates of cervical cancer are higher in Appalachian areas of Ohio, Kentucky, West

Virginia, and Virginia when compared to other parts of the U.S.⁸ Similarly, this area has lower screening and vaccination (for Human Papilloma Virus) rates, and high smoking rates. These rates are attributed to a region that is rural, often described as geographically isolated, with above-average poverty rates, low household income, and below-average educational attainment.⁹

Identified barriers to cervical cancer screening include:

- Preference for a female provider^{10,11}
- Competing priorities, such as caregiving and lack of childcare^{11,12}
- Lack of patient-centered communication: demeaning or discriminatory attitudes towards women (language, culture, low socio-economic status)¹¹
- Lack of accommodation for women's logistical needs around clinic hours and location¹¹
- Human papillomavirus infection (HPV) self-sampling-specific: forgetting, fear, lack of time, worry about using test incorrectly.¹³

Colorectal Cancer

Colorectal cancer is preventable with routine screenings. However, it remains the second leading cause of cancer mortality in the United States. In Appalachia, colorectal cancer incidence and mortality rates are higher when compared to the U.S.¹⁴ There are many factors attributable to the high mortality and incidence rates including unfamiliarity and lack of adherence to screening guidelines, feelings of worry about colorectal cancer, and poor overall health.

Identified barriers to colorectal cancer screening include:

- Lack of knowledge about stool-based testing options (e.g., FIT, FOBT) and appropriate frequency of each, doubt about the quality of tests, incorrect completion, inconvenience (e.g., some need multiple samples)¹⁵
- Colonoscopy-specific: perceived discomfort of prep and/or procedure, sense of violation (especially among men), embarrassment, and privacy and/or confidentiality concerns, especially in rural areas where people may personally know healthcare staff^{15,16}

 Some in primary care (especially pre-pandemic) recommended colonoscopy over home-based stool-testing, or only offered stool-based testing if colonoscopy was refused, instead of engaging in shared decision-making.¹⁵

Genetic Services

Generally, genetic services are more available around major cities, which creates a problem for those living in rural Appalachia. Tele-genetics, or the provision of remote genetics services, is a promising strategy for extending reach into rural areas and addressing distance-related access barriers. Further unintended access challenges may occur, however, for some rural, older, etc. patients in Appalachia due to lack of equipment or connectivity issues.¹⁷

Identified genetic services barriers include:

- Distance, transportation, and/or time barriers¹⁸⁻²⁰
- Technological access, literacy, and infrastructure limitations, such as lack of internet and mail service coverage^{5,18,21}
- High cost perceptions¹⁸
- Low prioritization^{19,21}
- Lack of physician recommendation (in part because of lack of physician skills and knowledge around genetic risk assessment, referral, and guidelines)^{18,21,23}
- Lack of insurance coverage and/or costs.^{18,21}

Interventions

Cervical Cancer

Numerous factors appear to facilitate cervical cancer screening. These factors include prior screening participation, referral by a healthcare professional, experience of positive symptoms, perceived need, prior HPV diagnosis, perception of convenience, interest in health status, and having family or friends who had cancer experiences.^{20,23,24}

Specific facilitating interventions and strategies include:

• HPV self-sampling

- Patient navigation and/or community health workers
- Community outreach
- Patient education
- Material supports
- Clinic-level interventions
- Community and/or systems-level interventions
- Policy interventions.

Colorectal Cancer

Colorectal cancer screening rates are affected by provider recommendation, family support and encouragement, perceived risk, family history of colorectal cancer, and knowing people with this type of cancer.^{11,13,20}

A systematic review found these interventions were associated with increased colorectal cancer screening completion: outreach, patient navigation, patient education, patient reminders, clinician interventions of academic detailing, clinician reminders, repeated mailed FOBT (fecal occult blood tests) with navigation.¹⁶

In fact, interventions designed to remove barriers for patients have increased colorectal cancer screenings by 37 percent.²⁵ Interventions specifically focused on patient navigation services provided through healthcare systems. Patient navigation services target populations experiencing greater disparities in cancer screening, including historically disadvantaged racial and ethnic populations, as well as those with lower incomes.

Specific facilitating interventions and strategies include:

- Test choice
- Community outreach
- Community outreach
- Patient education
- Material supports
- Clinic-level interventions
- Community and/or systems-level interventions
- Policy interventions.

Continuity of Care

As diagnosis and treatment options evolve in cancer care, so too do the number of providers patients see for their care. The cancer care continuum spans primary care through specialty and sub-specialty care providers. Collaboration between PCPs and specialty care providers can be challenging, and there are many opportunities to develop issues around care transitions. Continuity of care, therefore, becomes imperative to ensure quality outcomes for patients. The focus for continuity of care is around care transitions early in the cancer continuum, interactions between primary care and oncology, and follow-up after an abnormal cervical or colorectal screening.

Primary Care and Cancer Specialist Team Interactions

Primary care providers interact with specialty providers regularly. Those conversations and interactions are important regarding patient care. The landscape analysis found the following concerns among providers:

- Primary care providers prefer more communication from cancer specialists and report a significant gap between diagnosis through end of treatment. Oncologists also report infrequent communication and see room for improvement.²⁷
- Primary care providers prefer a shared-care model, oncologists prefer specialist-based care.²⁸

Cervical Cancer

There were logistical and psychological barriers identified for follow-up care after an abnormal (positive) Pap test result.²⁹

Colorectal Cancer

The main barrier under continuity of care for colorectal cancer was follow-up care after an abnormal (positive) FIT test result.³⁰ Secondary barriers included cost concerns, lack of insurance, lack of transportation, and psychological state after a positive screen.

Interventions

Strategies identified to mitigate care continuity issues and improve provider communication include:

• Using a memorandum of understanding (MOU) between primary care and local cancer programs

to formalize workflows and shared care processes

- Examining feasibility of shared electronic health records (EHR) or communication systems
- Agreeing to a referral system among providers
- Piloting embedding an oncologist in the primary care setting, or vice versa
- Examining opportunities for informal rapport-building between multidisciplinary care teams
- Identifying and sharing primary contact information for ease of ongoing contact
- Ensuring clear documentation including follow-up recommendations³¹
- Sharing data on cancer volume and stage to encourage collaboration among oncology programs
- Engaging with cancer care networks to collaborate on funding.

Interventions identified in the landscape analysis for cervical and colorectal cancer focus on provider-to-provider communications (examining how specialists such as cardiologists or endocrinologists interact with primary care providers), medical centers providing rural sites with technical assistance and services, and patient navigation.³²

Key Highlights from the Landscape Analysis

- Incidence rates for colorectal cancer in Appalachia range from 19.9 per 100,000 in Virginia to 29.5 in Mississippi (2015 to 2019) compared to the U.S. national rate of 21.8. Four Appalachian states (Mississippi, Kentucky, West Virginia, and Pennsylvania a) fall within the highest late-stage incidence, and five (Mississippi, West Virginia, Kentucky, Tennessee, Alabama) fall within the highest mortality quintile of states in the U.S.³¹
- Parts of Appalachia have been identified as early onset colorectal cancer hotspots.32 Colorectal cancer screening prevalence among people aged 50 and above ranged from 68.97 percent in Mississippi to 74.83 percent in Pennsylvania, compared to 71.9 percent in the U.S, with nine states above the national average.³¹
- Incidence rates for cervical cancer in Appalachia range from 2.6 in Virginia to 5.1 in Kentucky compared to the U.S. national rate of 3.6. Mortality rates range from 1.9 in Virginia, New York, and North Carolina, to 3.4 in Mississippi compared to the U.S. national rate of 2.2.³¹

Next Steps

The Alliance has chosen to focus its efforts on optimizing colorectal cancer screenings in Appalachia for 2023. The Alliance will prioritize relationship building among multidisciplinary providers through an e-newsletter, local, regional, and/or national meetings, and educational webinars. In addition, the Alliance will use an implementation framework to map evidence-based solutions to identified barriers and leverage existing resources and Alliance expertise to develop new solutions. Information and materials created will be disseminated amongst ACCC member programs and partners. Organizations or individuals interested in staying informed or participating in the Alliance can **sign up on the website accc-cancer.org/acca.**

For more information on the **Appalachian Community Cancer Alliance** and the Rural Appalachian Lung Cancer Screening Initiative, visit **ACCC's website accc-cancer.org/acca.**

References

- Appalachian Regional Commission. About the Appalachian region. Published January 4, 2023. Accessed January 18, 2023. https:// www.arc.gov/wp-content/uploads/2021/02/Health_Disparities_in_ Appalachia_Mortality_Domain.pdf.
- 2021-2022 Conference Series on Genetics Services in Appalachia. Accessed August 29, 2022. https://pharmacy.hsc.wvu.edu/ continuing-education/2021-2022-conference-series-on-genetics-services-in-appalachia/.

- Addressing cancer disparities in Appalachia. National Cancer Institute. Accessed January 20, 2023. https://www.cancer.gov/research/areas/ disparities/electra-paskett-addressing-cancer-disparities-appalachia.
- Morrow A, Chan P, Tucker KM, et al. The design, implementation, and effectiveness of intervention strategies aimed at improving genetic referral practices: a systematic review of the literature. Genet Med. 2021;23(12):2239-2249. doi:10.1038/s41436-021-01272-0
- Mikat-Stevens NA, Larson IA, Tarini BA. Primary-care providers' perceived barriers to integration of genetics services: a systematic review of the literature. Genet Med. 2015;17(3):169-176. doi:10.1038/ gim.2014.101
- Brady LA, Tumiel-Berhalter LM, Schad LA, et al. Increasing breast, cervical, and colorectal cancer screenings: A qualitative assessment of barriers and promoters in safety-net practices. J Pat Centered Res Rev. 2021; 8(4):323-330. doi: 10.17294/2330-0698.1857
- Fullenkamp AN, Haynes EN, Meloncon L, et al. Perceptions of genetic research in three rural Appalachian Ohio Communities. J Comm Genet. 2013; 4(1):9-17. doi: 10.1007/s12687-012-0112-2
- Weaver SJ, Jacobsen PB. Cancer care coordination: Opportunities for healthcare delivery research. Transl Behav Med. 2018; 8(3):503-508. doi: 10.1093/tbm/ibx079
- Paskett, ED. Covid-19 exacerbates cervical cancer disparities in Appalachia. Healio. Published March 4, 2021. Accessed January 19, 2023. https://www.healio.com/news/hematology-oncology/20210303/ covid19-exacerbates-cervical-cancer-disparities-in-appalachia
- Nalley C. Cervical cancer care: Focusing on disparities in Appalachia: Oncol Times. 2021;43(4):13,19. doi: 10.1097/01. COT.0000735036.34773.16
- Collins T, Stradtman LR, Vanderpool RC, et al. A community-academic partnership to increase Pap testing in Appalachian Kentucky. Am J Prev Med. Aug 2015;49(2):324-30. doi:10.1016/j.amepre.2015.04.028
- Majid U, Kandasamy S, Farrah K, et al. Women's preferences and experiences of cervical cancer screening in rural and remote areas: a systematic review and qualitative meta-synthesis. Rural Remote Health. 2019; 9(4):5190. doi: 10.22605/RRH5190
- Garcia C, Lothamer H, Mitchell EM. Provider-identified barriers to cervical cancer screening and perceptions toward self-collection of human Papillomavirus in Southwest Virginia. Public Health Nursing. 2016;33(6):539-546. doi:https://doi.org/10.1111/phn.12285
- Reiter PL, Shoben AB, McDonough D, et al. Results of a pilot study of a mail-based human Papillomavirus self-testing program for underscreened women from Appalachian Ohio. Sex Transm Dis. Mar 2019;46(3):185-190. doi:10.1097/olq.00000000000944
- 15. Oelsner WK, Shreenath AP, Sohail M, et al. An Emerging Health Disparity: Early Onset Colorectal Cancer in Appalachia. Conference: American College of Gastroenterology 2021 Meeting. Published October 21, 2021. Accessed April 12, 2023. https://www.researchgate. net/publication/355750306_An_Emerging_Health_Disparity_Early_ Onset_Colorectal_Cancer_in_Appalachia
- 16. Wang H, Roy S, Kim J, et al. Barriers of colorectal cancer

screening in rural USA: a systematic review. Rural Remote Health. Aug 2019;19(3):5181. doi:10.22605/rrh5181

- Hatcher J, Dignan MB, Schoenberg N. How do rural health care providers and patients view barriers to colorectal cancer ccreening? Insights from Appalachian Kentucky. Nursing Clinics. 2011;46(2):181-192. doi:10.1016/j.cnur.2011.02.001
- Terry A. NYMAC experiences with genetics services in Appalachia email correspondence 9/19/22. In: Phillips S, editor.
- West Virginia University, School of Pharmacy. (2022). 2021-2022 Conference Series on Genetics Services in Appalachia. Accessed August 29, 2022. https://pharmacy.hsc.wvu.edu/continuing-education/2021-2022-conference-series-on-genetics-services-in-appalachia/
- Morrow A, Chan P, Tucker KM, et al. The design, implementation, and effectiveness of intervention strategies aimed at improving genetic referral practices: a systematic review of the literature. Genet Med. 2021;23(12):2239-2249. doi:10.1038/s41436-021-01272-0
- Crosby RA, Stradtman L, Collins T, et al. Community-based colorectal cancer screening in a rural population: who returns fecal immunochemical test (FIT) kits? J Rural Health. 2017;33(4):371-374. doi:https://doi. org/10.1111/jrh.12210
- 22. Institute of Medicine (US) and National Research Council (US) National Cancer Policy Board; Curry SJ, Byers T, Hewitt M, editors. Fulfilling the Potential of Cancer Prevention and Early Detection. Washington (DC): National Academies Press (US); 2003. 5, Potential of Screening to Reduce the Burden of Cancer.
- Byrne L, Morman N, Shane-Carson KP, et al. Genetic counseling referrals among cancer registry patients who meet NCCN guidelines: an Ohio Study. Oncol Issues. 2022;37(4).
- Institute of Medicine (US) and National Research Council (US) National Cancer Policy Board; Curry SJ, Byers T, Hewitt M, editors. Fulfilling the Potential of Cancer Prevention and Early Detection. Washington (DC): National Academies Press (US); 2003. 5, Potential of Screening to Reduce the Burden of Cancer.
- Hall MB, Vos P, Bess JJ, et al. Cervical cancer screening behaviors and perceptions of medical mistrust among rural Black and White women. J Health Care Poor Underserved. 2018;29(4):1368-1385. doi:10.1353/ hpu.2018.0101
- Tsui J, Howard J, O'Malley D, et al. Understanding primary care-oncology relationships within a changing healthcare environment. BMC Family Practice. 2019;20(1):164. doi:10.1186/s12875-019-1056-y
- U.S. Centers for Disease Control and Prevention. Reducing structural barriers. Published December 1, 2022. Accessed January 27, 2023. https://www.cdc.gov/screenoutcancer/interventions/reducing-structural-barriers.htm.
- Schoenberg NE, Howell BM, Fields N. Community strategies to address cancer disparities in Appalachian Kentucky. Fam Community Health. 2012; 35(1):31-43. doi: 10.1097/FCH.0b013e3182385d2c
- Dossett LA, Hudson JN, Morris AM, et al. The primary care provider (PCP)-cancer specialist relationship: A systematic review and mixed-methods meta-synthesis. CA Cancer J Clin. 2017;67(2):156-169.

doi:https://doi.org/10.3322/caac.21385

- Ely GE, White C, Jones K, et al. Cervical cancer screening: exploring Appalachian patients' barriers to follow-up care. Soc Work Health Care. 2014;53(2):83-95. doi:10.1080/00981389.2013.827149
- Falah N, Mallow J, Kelly K. Telegenetics in Rural Appalachia: Implementation Experiences and Barriers to Effective Clinical Care for Populations Experiencing Health Disparity. Published March 28, 2022. Accessed April 12, 2023. https://assets.researchsquare. com/files/rs-1407787/v1/c402daa5-d4ed-466c-ad6e-6f23bd696efe. pdf?c=1650278665
- Kentucky Cabinet for Health and Family Services. Colon Cancer Prevention Program. Accessed September 20, 2022. https://chfs.ky.gov/ agencies/dph/dpqi/cdpb/Pages/coloncancer.aspx
- Malo TL, Correa SY, Moore AA, et al. Centralized colorectal cancer screening outreach and patient navigation for vulnerable populations in North Carolina: study protocol for the SCORE randomized controlled trial. Implement Sci Commun. 2021;2(1):113. doi:10.1186/ s43058-021-00194-x

Acknowledgements

Thank you to Bristol Myers Squibb for their support of this program.



Association of Community Cancer Centers

The Association of Community Cancer Centers (ACCC) is the leading education and advocacy organization for the cancer care community. Founded in 1974, ACCC is a powerful network of 30,000 multidisciplinary practitioners from 2,000 hospitals and practices nationwide. As advances in cancer screening and diagnosis, treatment options, and care delivery models continue to evolve—so has ACCC—adapting its resources to meet the changing needs of the entire oncology care team. For more information, visit **accc-cancer.org**.

 $\hfill {\ensuremath{\mathbb O}}$ 2023. Association of Community Cancer Centers. All rights reserved. No part of this publication may be reproduced or transmitted in any form or by any means without written permission.

This publication is a benefit of ACCC membership.