Genentech | ACCESS SOLUTIONS®
A Member of the Roche Group

for TECENTRIQ® (atezolizumab)

SAMPLE CODING

Hepatocellular Carcinoma (HCC)

TYPE	CODE		DESCRIPTION
Diagnosis: ICD-10-CM	C22.0		Liver cell carcinoma, hepatocellular carcinoma
	C22.8		Malignant neoplasm of liver, primary, unspecified as to type
Drug: HCPCS	J9022		Injection, atezolizumab, 10 mg
Drug: NDC Note: Payer requirements regarding use of a 10-digit or 11-digit NDC may vary. Both formats are listed here for your reference.	10-digit	11-digit	
	50242-917-01	50242-0917-01	1200 mg/20 mL single-dose vial
	50242-918-01	50242-0918-01	840 mg/14 mL single-dose vial
Administration procedures: CPT	96413		Chemotherapy administration, intravenous infusion technique; up to 1 hour, single or initial substance/drug
	96415		Chemotherapy administration, intravenous infusion technique; each additional hour (List separately in addition to code for primary procedure)
	96417		Chemotherapy administration, intravenous infusion technique; each additional sequential infusion (different substance/drug), up to 1 hour (List separately in addition to code for primary procedure)

CPT=Current Procedural Terminology; HCPCS=Healthcare Common Procedure Coding System; ICD-10-CM=International Classification of Diseases, 10th Revision, Clinical Modification; NDC=National Drug Code.

These codes are not all-inclusive; appropriate codes can vary by patient, setting of care and payer. Correct coding is the responsibility of the provider submitting the claim for the item or service. Please check with the payer to verify codes and special billing requirements. Genentech does not make any representation or guarantee concerning reimbursement or coverage for any service or item.

Many payers will not accept unspecified codes. If you use an unspecified code, please check with your payer.

INDICATIONS & IMPORTANT SAFETY INFORMATION

INDICATIONS

HCC

TECENTRIQ, in combination with bevacizumab, is indicated for the treatment of patients with unresectable or metastatic hepatocellular carcinoma (HCC) who have not received prior systemic therapy.

IMPORTANT SAFETY INFORMATION

Serious Adverse Reactions

Please refer to the full Prescribing Information for important dose management information specific to adverse reactions.

- **Immune-mediated pneumonitis.** Immune-mediated pneumonitis or interstitial lung disease, including fatal cases, have occurred. Permanently discontinue TECENTRIQ for Grade 3 or 4 pneumonitis

IMPORTANT SAFETY INFORMATION (cont)

- **Immune-mediated hepatitis.** Immune-mediated hepatitis and liver test abnormalities, including fatal cases, have occurred. Permanently discontinue TECENTRIQ for AST or ALT elevations more than 8 times the upper limit of normal or total bilirubin more than 3 times the upper limit of normal
- **Immune-mediated colitis.** Immune-mediated diarrhea or colitis have occurred. Permanently discontinue TECENTRIQ for Grade 4 diarrhea or colitis
- **Immune-mediated endocrinopathies.** Thyroid disorders, adrenal insufficiency, type 1 diabetes mellitus, including diabetic ketoacidosis, and hypophysitis/hypopituitarism have occurred
- **Other immune-mediated adverse reactions.** TECENTRIQ can cause severe and fatal immune-mediated adverse reactions. These immune-mediated reactions may involve any organ system. Permanently discontinue TECENTRIQ for Grade 4 immune-mediated adverse reactions involving a major organ
- **Infections.** Severe infections, including fatal cases, have occurred
- **Infusion-related reactions.** Severe or life-threatening infusion-related reactions have occurred. Permanently discontinue TECENTRIQ in patients with Grade 3 or 4 infusion-related reactions
- **Embryo-fetal toxicity.** TECENTRIQ can cause fetal harm in pregnant women. Verify pregnancy status prior to initiating TECENTRIQ. Advise females of reproductive potential of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TECENTRIQ and for at least 5 months after the last dose
- Advise female patients not to breastfeed while taking TECENTRIQ and for at least 5 months after the last dose

Most Common Adverse Reactions

The most common adverse reactions (rate ≥20%) in patients who received TECENTRIQ in combination with bevacizumab for HCC were hypertension (30%), fatigue/asthenia (26%), and proteinuria (20%).

You may report side effects to the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. You may also report side effects to Genentech at 1- 888-835-2555.

Please see full Prescribing Information for additional Important Safety Information.

TECENTRIQ®, its logo and the Access Solutions logo are registered trademarks of Genentech, Inc.

Please see full Prescribing Information for additional Important Safety Information.

Genentech
A Member of the Roche Group
©2020 Genentech USA, Inc. So. San Francisco, CA
All rights reserved. M-US-00004118(v1.0) 06/20